





# **SOHO** – The Solar and Heliospheric Observatory




**SOHO is a mission of  
international cooperation  
between ESA and NASA**



**Artist's rendition of the SOHO spacecraft**





**SOHO spacecraft being prepared for thermal tests  
at Intespace in Toulouse, France**





**SOHO payload module, without thermal blankets,  
at the end of its integration and testing at  
Matra Marconi Space**





# The SOHO Spacecraft

1 **SUMER**: Solar Ultraviolet Measurements of Emitted Radiation

2 **CDS**: Coronal Diagnostic Spectrometer

3 **EIT**: Extreme-ultraviolet Imaging Telescope

4 **UVCS**: UltraViolet Coronagraph Spectrometer

5 **LASCO**: Large-Angle and Spectrometric Coronagraph

6 **SWAN**: Solar Wind ANisotropies

11 **MDI**: Michelson Doppler Imager


10 **VIRGO**: Variability of solar Irradiance and Gravity Oscillations

9 **GOLF**: Global Oscillations at Low Frequencies


8 **COSTEP**: Comprehensive SupraThermal and Energetic Particle analyser

7 **CELIAS**: Charge, Element and Isotope Analysis System

8 **ERNE**: Energetic and Relativistic Nuclei and Electron experiment




SOHO, a solar scientific observatory, has 12 instruments on board to observe the Sun 24 hours a day. It is a mission of international cooperation between ESA and NASA.




**Schematic of SOHO's orbital path in relation to the Earth, moon, and Sun – SOHO is about 1.5M km sunward of the Earth**






Ground operations lost contact with SOHO on 24 June 1998, but through diligent efforts the recovery team was able to nurse SOHO back to life by November 1998




**A composite image of the Sun that depicts the range of SOHO's scientific research from the solar interior, to the surface and corona, and out to the solar wind**






**The parts of the Sun**




**Erupting prominence as recorded by EIT  
in the He II 304Å line**






**Active regions and magnetic loops as recorded  
by EIT in the Fe IX/X 171Å line**





**EIT composite image from three wavelengths (171Å, 195Å and 284Å) revealing solar features unique to each wavelength**





**Ratio of EIT full Sun images in  
Fe XII 195Å to Fe IX/X 171Å – Bright  
areas are hotter; dark areas are cooler**





**Magnetic loops and prominences captured by the Extreme ultraviolet Imaging Telescope (EIT) in three wavelengths**






EIT 171 full disk image



25 April 1998


**Close-up of an active region in extreme ultraviolet light  
from NASA's TRACE (Transition Region and Coronal  
Explorer) spacecraft**




**A series of EIT 195Å images over two days shows two active regions connecting their magnetic field lines over a large area of the Sun**

**Images are Fe XII at 195Å showing the solar corona at a temperature of about 1.5 million K.**







Large, eruptive prominence in He II at 304Å, with an image of the Earth added for size comparison



**26 August 1997**




**One of the largest eruptive prominences recorded by SOHO/EIT in 1997 in He II at 304Å. It reached 28 times the size of Earth.**







Ne V - 400,000 K




O V - 250,000 K




O III - 85,000 K



He II - 50,000 K




He I - 20,000 K




Monochromatic images  
obtained simultaneously from  
CDS representing different  
temperatures and structures  
of the solar atmosphere


Ca X - 630,000 K




Mg IX - 1,000,000 K




Fe XII - 1,600,000 K




Fe XIV - 2,000,000 K



Fe XVI - 2,600,000 K






**SUMER image in S VI at 933 Å  
on 12 May 1996**






**Iron XII at 1242.01Å**  
(corona)  
1.6 million K



**Magnesium X at 624.943Å**  
(corona)  
1.1 million K




**Oxygen V at 629.729Å**  
(transition region)  
230,000 K




**Nitrogen V at 1238.82Å**  
(transition region)  
180,000 K



**UV – Continuum emission at 1240Å**  
(chromosphere)  
10,000 K






**SUMER scans of a north polar coronal hole in lines  
formed at temperatures from 10,000 to 1.6 MK**






EIT 195Å

Oxygen VI



Lyman beta





Intensity

Velocity

Line width

**Simultaneous imaging of UV emission, gas flow velocities, and spectral line width of active region loop structures observed with SUMER**






**Supersonic flow velocities observed by CDS during a solar eruption from the south west limb. The velocities approach 300 km/s at the leading edge (B).**




## Flows in an Active Region Loop System

EIT Fe XII 195 Å



July 27 1997 22:44 UT


O V 629 Å



MONOCHROMATIC IMAGE FROM  
CORONAL DIAGNOSTIC SPECTROMETER (CDS)

July 27, 1996

CDS LINE PROFILES




Active region loop system above the east limb observed in O V on 27 July 1996 by CDS. The line profiles from three different spatial locations (A, B, and C) are displayed in the right panel.







EIT 304 full disk image



Velocity image CDS, OV (250,000° K)



Solar “tornado” observed by SOHO/CDS with speeds up to 500,000 km/h




**SUMER Doppler velocity map (close-up) of a polar coronal hole region showing the source regions of the fast solar wind. The strongest flows occur near the boundary intersections of the supergranular network cells.**






UVCS

SUMER




Line profile of O VI from UVCS observed in a polar coronal hole (lower panel) compared to disk observation from SUMER. The extremely broad O VI line yields velocities up to 500 km/s, which corresponds to a kinetic temperature of 200 million K.




**UVCS/SOHO has discovered surprisingly fast spiraling motions for charged oxygen atoms (1.3 million Km/h) over the solar poles, as compared to hydrogen atoms (at about half this speed). The coiling speeds of oxygen atoms along the spirals are 20 times larger than were expected.**





**MDI Full Disk Magnetogram  
9 May 1999**




**MDI magnetogram overlay with lines of convergence of the horizontal flow. Green dots show the convergence points. Measured flow is shown as colored arrows (red= downflow; blue= upflow).**







CDS OV intensity




MDI magnetogram



Overlay of CDS and MDI



**Correlation of transition region EUV emission and  
photospheric magnetic fields from observations by  
CDS and MDI**







**Model of magnetic field lines above the solar surface based on MDI magnetic field measurements superimposed on EIT coronal emission observations**





5 November 1998




*Close-up magnetogram image of sunspots*


**An EIT 304Å image, an MDI full disk white light image, with a close-up, and a high-resolution magnetogram all view the same magnetic structures that we call sunspots.**



**White light  
image**




**Magnetogram**



**Dopplergram**




**$k$ - $\omega$  p mode  
spectrum**




**SOHO/MDI high resolution images of an active  
region, taken on 5 November 1998 after successfully  
recomissioning of the instrument**






**The subsurface structure (sound speed) below a sunspot  
as derived from Doppler measurements by MDI**




**A bright solar flare captured on 2 May 1998 in the  
195Å line of Fe XII**

**(The horizontal white line on either side of the flare was caused by  
charge bleeding on the CCD detector.)**






**A rapidly expanding “solar quake” on the Sun’s surface based on data from the Michelson Doppler Imager (MDI). The wave was caused by a solar flare on 6 July 1996 and spread out more than 100,000 km at the solar surface.**




**EIT observation of a shock wave expanding across much of the Sun's surface from a coronal mass ejection (CME) initiation site on 12 May 1997. This “running difference” imaging technique emphasizes the changes between successive frames.**






**An image sequence showing the progress over eight hours of a clearly defined coronal mass ejection on 5-6 August 1999 taken by LASCO C3.**




**LASCO C3 image of the large coronal mass ejection  
(CME) of 20 April 1998**






**A composite of four images of a large CME from 6 November 1997, which was associated with an X-9.4 flare**



**LASCO C2 coronagraph image in which a twisting, helical-shaped CME spins off from the Sun**













**A LASCO C2 “running difference” image showing a “halo”  
CME blast beginning its journey towards Earth**




# The Solar Cycle



The rise of activity cycle 23 as reflected by the number of sunspots recorded to date and as projected (dotted line). Selected EIT 195Å images and MDI magnetograms are shown.






**The Sun's magnetic field and plasma releases directly affect Earth and the rest of the solar system. This schematic view illustrates a magnetic storm approaching Earth and how the solar wind shapes the Earth's magnetosphere.**




## Normal magnetosphere




## Magnetosphere being affected by a CME

When the particles from a CME impact the Earth's magnetosphere, the sunward side flattens and the tail elongates. Note that most particles are drawn in on the far side.







**The numerous effects of space weather**



Credit: Jan Curtis




**An aurora, the most spectacular visual effect  
of magnetic storms seen on Earth**





## Single Dopplergram

(30-MAR-96 19:54:00)



SOI / MDI

Stanford Lockheed Institute for Space Research


## MDI Full Disk Dopplergram

(dark colors = motion toward the observer)



## Average Dopplergram Minus Polynomial Fit

45 images averaged (30-Mar-96 19:26 to 30-Mar-96 20:17)



SOI / MDI


Stanford Lockheed Institute for Space Research

**MDI Full Disk Dopplergram**  
showing superanular convection cells  
on the Sun's surface






Single Dopplergram Minus 45 Images Average  
(30-MAR-96 19:54:00)



SOI / MDI

Stanford Lockheed Institute for Space Research

**MDI Full Disk Dopplergram**  
showing the p-mode oscillations of the Sun




**Power spectrum obtained from 2 months of continuous MDI data (May/June 1996). The "ridges" of greater power result from standing sound waves resonating within the Sun.**






## Structure of the Solar Interior




**Fourier spectrum of global oscillations  
observed by GOLF**



**Temporal variations of the amplitudes of solar p-modes as measured by VIRGO**






**An MDI dopplergram image of the Sun's surface is merged with a helioseismology image of the flows of plasma in the solar interior. The smaller diagram shows the paths of several different acoustic (pressure) waves inside of the Sun whose measurements reveal its internal structure.**




Jet streams




**Polar plasma jet streams and variation in rotational speed in the solar interior as measured by MDI**






**Solar rotation and polar flows of the Sun as deduced from measurements by MDI. The cutaway reveals rotation speed inside the Sun.**




**Radial and latitudinal variations of the sound speed in the Sun as derived from MDI and VIRGO measurements. Red = hotter regions than in standard model, blue = cooler regions.**






**First image of the sub-surface temperatures and flows in the convection zone of a star deduced from MDI observations**




**CELIAS chart showing new (*red*) and rarely observed (*green*) elements and isotopes discovered in the solar wind**






**Series of solar energetic particle events observed  
in July 1996 by the COSTEP instrument**




**Time profile of elements during the flare event of 6 November 1997, as detected by the ERNE instrument**





**Galactic cosmic rays as recorded by ERNE. Galactic cosmic radiation consists of particles originating in the Milky Way**






**Lyman- $\alpha$  whole sky map as recorded by SWAN on 2 February 1996. The U-shaped yellow, bright band is the Milky Way.**




**SWAN observation of active regions on the far side of the Sun.  
Active regions illuminate the distant interstellar hydrogen cloud  
like a searchlight strikes clouds at night.**




**Comet Hale-Bopp seen approaching the Sun in a time series of six SWAN full sky images in the ultraviolet light (1100-1800 Å) from 4 January to 3 April 1997**






**SWAN recording of the huge cloud of hydrogen, 70 times the size of the Sun, surrounding Comet Hale-Bopp when it neared the Sun in 1997. Ultraviolet light revealed a cloud 100 million kilometres wide and diminishing in intensity outwards (contour lines).**




## Observation of the shadow of Comet Hale-Bopp by SWAN




2 June 1998

**Two “Sungrazing” comets heading in tandem towards the Sun’s corona. They do not reappear on the other side.**





**A comet observed by UVCS in Lyman alpha on  
1-2 May 1997 as it approaches the Sun**





**Spectacular view of the solar corona with Venus and a sungrazing comet (SOHO 58) as observed by LASCO**





*LASCO C2*

*LASCO C3*



1997



1998



1999




The changing shape and structure of the corona  
with the solar cycle






**1997 January 23**








**1998 November 9**




**A comparison of two EIT images almost two years apart illustrates how the level of solar activity has increased significantly**

**Images are Fe IX/X at 171 Å showing the solar corona at a temperature of about 1 million K.**




**Increasing total solar irradiance as measured by VIRGO since SOHO's launch. The EIT full disk images show a corresponding increase in solar activity.**



**Total irradiance variations during solar cycles 21–23 as recorded by several satellites since 1978. The data shaded in green is from the VIRGO instrument.**





**The gradual increase in solar activity as shown in the EIT and LASCO C3 images illustrates the approach of solar maximum**